



2019 CONJOINT CONFERENCE CONFÉRENCE CONJOINTE DE 2019

Admissions due to Primary Care Sensitive Conditions: is there an association with the acquisition of Healthcare-Associated Infection?

Ercília Evangelista Porto, Thaís Guimarães, Maria Clara Padoveze

School of Nursing, University of São Paulo

Faculty of Medicine, University of São Paulo


Brazil

Study funded by the National Council for Scientific Technological Development (CNPq), process 460783/2014-9_

Background

Brazil

- ❑ Healthcare-associated infection (HAI) is a major public health problem.
- ❑ Primary Care:
 - ❑ Key component of healthcare
 - ❑ Integrating element of the healthcare system.
 - ❑ Potential to prevent unnecessary hospital thus indirectly prevents exposing the patient to iatrogenic risks, such as HAI.
- ❑ Primary Care Sensitive Conditions (PCSC): indicator that measures the effectiveness of primary care.



Objective

To identify whether there is an association between PCSC and HAI.

- Hypothesis: are PCSC patients admitted more susceptible to HAI?
- Literature: no previous study investigating the association of PCSC with HAI



São Paulo. Source: google images

Clinics Hospital, USP. Source: google images

Methods



- ❑ Study design: prospective cohort.
- ❑ Setting: Clinics Hospital, tertiary and reference in São Paulo.
 - ❑ 910 beds (107 beds for intensive care)
- ❑ Annually: 26,000 elective surgeries; 37,000 consultations (high complexity patients)
- ❑ from May 2018 to December 2018
- ❑ Sample: 605 inpatients
- ❑ This project was approved by our institutional Ethics Committee

Methods

Data Collection Sources

- Patients' interviews: structured questionnaire to characterize the potential social determinants of the participants.
- Patients' medical records: PCSC and other variables
- Identification of HAI cases: as per routine of infection control team

Criteria

Inclusion: more than 48 hours of hospitalization

HAI:

- Brazilian standardized criteria, adapted from CDC, EUA.

PCSC:

- Brazilian national list

Analysis: descriptive statistic and univariate analysis; significance of <0.05 .

Results

- Patients were 57% females (n=345)
- The majority of patients were black or brown (n= 524; 86.6%)
- Average age: 52.8 years
- Average schooling: 8.5 years
- Average family income: 2.5 national minimum wage;

- The Human Development Index (HDI) of origin: varied from 0.72 to 0.81
- Gini Index varied from 0.41 to 0.67.

Figure 1. Distribution of the origin of participants. São Paulo, 2018



Results

Patients with PCSC and HAI: OR 1.93
(95% IC -0.714-5.252), p=0.199.

Risk Difference: 4.2

Table 1. Distribution of HAI and PCSC. São Paulo, 2018.

Variable	No HAI	With HAI
	N(%)	N(%)
No PCSC	523 (91.3)	27 (84.4)
With PCSC	50 (8.7)	5 (15.6)
Total	573 (100%)	32 (100%)

HAI n=32 (5.3%)

Most frequent:

Surgical Site: n=7 (21.8%)

Catheter-related bloodstream: n= 7 (21.8%)

Catheter-related urinary tract : n= 6 (18.8%)

PCSC n=55 (9.1%)

Most frequent:

Cerebrovascular disease: n= 17 (30.9%)

Diabetes mellitus: n= 10 (18.2%)

Kidney & urinary tract infection: n= 8 (14.5%)

Results

Table 2. Characteristics of patients with HAI & PCSC. São Paulo, 2018.

Patient	CID 10	PCSC	Other commorbidities	HAI	Etiologic agent
1 F, 69	N39	Urinary tract infection, unspecified	Systemic lupus erythematosus Arthritis Non-dialytic renal disease Diabetes mellitus	Osteomyelitis	<i>A. baumannii</i> resistant to carbapanems
2 M, 79	E 10.5	Insulin-dependant diabetes mellitus + peripheral circulatory complications	None	Osteomyelitis	<i>S. aureus</i> methicillin resistant
3 M, 80	E10.5	Insulin-dependant diabetes mellitus + peripheral circulatory complications	None	Osteomyelitis	<i>S. aureus</i> methicillin resistant
4 M, 50	J15.9	Bacterial pneumonia, unspecified	Diabetes mellitus	Cateter-related Bloodstream infection	<i>Salmonella spp</i>
5 M, 70	E43	Severe protein-calorie malnutrition, unspecified	None	Catheter-related Urinary tract	<i>C. freundii</i> complex

Results

No relevant differences among selected individual or social indicators when comparing patients with or without HAI

Table 3. Characteristics of patients with HAI or without HAI. São Paulo, 2018.

Variable	With HAI	Without HAI
Age, years (mean)	57	52.6
Charlson comorbidity index comorbidity	2.7	2.2
Schooling, years (mean)	8.8	8.5
Monthly wage (average)	R\$2,507	R\$2,751
HDI	0.74-0.86	0.71-0.88
Gini index	0.45-0.67	0.41-0.67

HDI: Human Development Index

Discussion

- Paker et al (2015): no overall association between Scottish Index of Multiple Deprivation and HAI.
- Fisman et al (2014): regions with lower gross domestic product dedicated to healthcare and proximity to Equator with higher probability of bloodstream infections caused by Gram-negatives
- Nomamiukor et al (2015): living conditions associated with increased antibiotic resistance in community isolates of *E. coli*.

Fisman et al. Plos One 2014; 9(12)

Nomamiukor et al. J Antimicrob Chemother 2015; 70(11):3154-8.

Paker et al. Journal of Hospital Infections 2015; 91: 351-357

Conclusions



- ❑ No statistical association between PCSC and HAI
- ❑ The overall risk difference pointed out that other studies are necessary to evaluate our hypothesis.
- ❑ Next step: a multilevel analysis will be performed to eliminate potential confounders that may abstruse the deep understanding of this phenomenon.

Thank you!

padoveze@usp.br